

Decoupling of heavy-quark loops in light-light and heavy-light quark currents

A. G. Grozin*

*Institute of Theoretical Particle Physics, University of Karlsruhe,
D-76128 Karlsruhe, Germany*

Abstract

Matching of light-light quark currents in QCD with a heavy flavour and in the low-energy effective QCD is calculated in $\overline{\text{MS}}$ at two loops. Heavy-light HQET currents are similarly considered.

Consider QCD with n_l light flavours and a single heavy one having mass M . Processes involving only light fields with momenta $p_i \ll M$ can be described by an effective field theory — QCD with n_l flavours plus $1/M^n$ suppressed local operators in the Lagrangian, which are the remnants of heavy quark loops shrunk to a point. This low-energy effective theory is constructed to reproduce S -matrix elements of the full theory expanded to some order in p_i/M . This condition fixes $\overline{\text{MS}}$ coupling $\alpha'_s(\mu)$ of the effective theory, its gauge parameter $a'(\mu)$ and running light quark masses $m'_i(\mu)$ in terms of the parameters of the full theory. Two-loop matching was considered in [1, 2], and three-loop one — in [3].

Operators of the full QCD can be expanded in $1/M$ in operators of the effective theory. Coefficients of these expansions are fixed by equating on-shell matrix elements. In this Letter, I consider bilinear quark currents $j_n(\mu) = Z_n^{-1}(\mu)j_{n0}$, where the bare currents are $j_{n0} = \bar{q}_0 \gamma^{[\mu_1} \dots \gamma^{\mu_n]} q_0$, and $Z_n(\mu)$ are their $\overline{\text{MS}}$ renormalization constants. In the low-energy theory, they are equal to $C_n(\mu)j'_n(\mu)$ plus power-suppressed terms (which will be omitted throughout this paper). It is natural to match the currents at $\mu = M$, because $C_n(M)$ contains no large logarithms. Currents at arbitrary normalization scales can be related by

$$\exp \left(- \int_{\alpha_s(M)}^{\alpha_s(\mu)} \frac{\gamma_n(\alpha)}{2\beta(\alpha)} \frac{d\alpha}{\alpha} \right) j_n(\mu) = C_n(M) \exp \left(- \int_{\alpha'_s(M)}^{\alpha'_s(\mu')} \frac{\gamma'_n(\alpha)}{2\beta'(\alpha)} \frac{d\alpha}{\alpha} \right) j'_n(\mu'), \quad (1)$$

where

$$\gamma_n(\alpha_s) = \frac{d \log Z_n}{d \log \mu} = \gamma_{n0} \frac{\alpha_s}{4\pi} + \gamma_{n1} \left(\frac{\alpha_s}{4\pi} \right)^2 + \dots \quad (2)$$

is the current's anomalous dimension (obtained at two loops for generic n in [4]), and

$$\frac{d \log \alpha_s}{d \log \mu} = -2\varepsilon - 2\beta(\alpha_s), \quad \beta(\alpha_s) = \beta_0 \frac{\alpha_s}{4\pi} + \beta_1 \left(\frac{\alpha_s}{4\pi} \right)^2 + \dots, \quad \beta_0 = \frac{11}{3}C_A - \frac{4}{3}T_F n_f, \quad (3)$$

*Permanent address: Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia. Email: A.G.Grozin@inp.nsk.su

$T_F = \frac{1}{2}$, $C_A = N$, $C_F = (N^2 - 1)/(2N)$, $N = 3$ is the number of colours; $n_f = n_l + 1$; $d = 4 - 2\varepsilon$ is the space-time dimension; γ'_n, β' are the corresponding quantities in n_l -flavour QCD.

On-shell matrix element of $j_n(\mu)$ is $M(\mu) = Z_q Z_n^{-1}(\mu) \Gamma_{n0}$, where Γ_{n0} is the bare vertex function and Z_q is the on-shell renormalization constant of the light-quark field. It should be equal to $C_n(\mu) M'_n(\mu)$ plus power-suppressed terms, where $M'_n(\mu) = Z'_q Z_n'^{-1}(\mu) \Gamma'_{n0}$. Both matrix elements are ultraviolet-finite; their infrared divergences coincide, because both theories are identical in the infrared region. Therefore (cf. [5, 4]),

$$C_n(\mu) = \frac{Z_q}{Z'_q} \frac{Z'_n(\mu)}{Z_n(\mu)} \frac{\Gamma_{n0}}{\Gamma'_{n0}}. \quad (4)$$

The on-shell wave-function renormalization constant is $Z_q = [1 - \Sigma_V(0)]^{-1}$, where the bare mass operator is $\Sigma(p) = \not{p} \Sigma_V(p^2)$. Only the heavy-quark loop contributes (Fig. 1a), and we obtain at two loops

$$\Sigma_V(0) = C_F T_F \frac{g_0^4 M^{-4\varepsilon}}{(4\pi)^d} I \frac{(d-1)(d-4)}{d}, \quad (5)$$

where

$$i C_F g_0^2 \int \frac{d^d k}{(2\pi)^d} \frac{\Pi(k^2)}{k^4} = C_F T_F \frac{g_0^4 M^{-4\varepsilon}}{(4\pi)^d} I, \quad (6)$$

$$I = \Gamma^2(\varepsilon) \frac{2(d-6)}{(d-2)(d-5)(d-7)} = -\frac{2}{3} \Gamma^2(\varepsilon) \left(1 - \frac{2}{3}\varepsilon + \frac{22}{9}\varepsilon^2 + \dots\right),$$

and $\Pi(k^2)$ is the heavy-quark contribution to the gluon polarization operator. This agrees with the decoupling relation [3] $q = (Z_q/Z'_q)^{1/2} q'$.




Figure 1: Mass operator and vertex function

The bare vertex function (Fig. 1b) can be calculated at any on-shell momenta of the quarks. It is most easy to set both momenta to zero (thus excluding power-suppressed terms in (4)):

$$\Gamma_{n0} = 1 - C_F T_F \frac{g_0^4 M^{-4\varepsilon}}{(4\pi)^d} I \left(\frac{(d-2n)^2}{d} - 1 \right). \quad (7)$$

The corresponding quantities in the low-energy theory are $Z'_q = 1$, $\Gamma'_{n0} = 1$. The ratio

$$\frac{Z_n}{Z'_n} = 1 + \frac{1}{4} (\gamma_{n0} \Delta \beta_0 - \Delta \gamma_{n1} \varepsilon) \left(\frac{\alpha_s}{4\pi \varepsilon} \right)^2 = 1 + \frac{1}{9} C_F T_F \left(\frac{\alpha_s}{4\pi \varepsilon} \right)^2 (n-1) [6(n-3) - (n-15)\varepsilon] \quad (8)$$

(where $\Delta\beta_0 = -\frac{4}{3}T_F$ and $\Delta\gamma_{n1}$ are the contributions of a single flavour) can be obtained from [4].

Using

$$\frac{g_0^2 M^{-2\varepsilon} \Gamma(1+\varepsilon)}{(4\pi)^{d/2}} = \frac{\alpha_s(M)}{4\pi} + \mathcal{O}(\alpha_s^2), \quad (9)$$

we finally obtain

$$j_n(M) = \left[1 + C_F T_F \left(\frac{\alpha_s}{4\pi} \right)^2 \frac{(n-1)(85n-267)}{54} \right] j'_n(M). \quad (10)$$

To all orders, $j_1 = j'_1$ [5]; this is natural, because the integral of the vector current is the number of quarks minus antiquarks, which is the same in both theories. Also $m(\bar{q}q) = m'(\bar{q}q)'$ to all orders [6]. Therefore, C_0 can be obtained from the mass decoupling relation [1, 3]. The currents j_3 and j_4 differ from j_1 and j_0 by insertion of 't Hooft-Veltman γ_5^{HV} ; they differ from the corresponding currents with the anticommuting γ_5^{AC} by finite renormalizations $\bar{q}\gamma_5^{\text{AC}}\gamma^\mu q = Z_A \bar{q}\gamma_5^{\text{HV}}\gamma^\mu q$, $\bar{q}\gamma_5^{\text{AC}}q = Z_P \bar{q}\gamma_5^{\text{HV}}q$, known to three loops [7]:

$$\begin{aligned} Z_A &= 1 - 4C_F \frac{\alpha_s}{4\pi} + C_F \left(\frac{\alpha_s}{4\pi} \right)^2 \frac{198C_F - 107C_A + 4T_F n_f}{9} + \dots, \\ Z_P &= 1 - 8C_F \frac{\alpha_s}{4\pi} + C_F \left(\frac{\alpha_s}{4\pi} \right)^2 \frac{2(C_A + 4T_F n_f)}{9} + \dots \end{aligned} \quad (11)$$

Insertion of γ_5^{AC} does not change the matching coefficient. Hence, $C_3 = C_1 Z'_A / Z_A$, $C_4 = C_0 Z'_P / Z_P$, where Z'_A , Z'_P contain n_l instead of $n_f = n_l + 1$. The result for $n = 2$ is, to the best of my knowledge, new.

The HQET heavy-light current $\tilde{j}(\mu) = \tilde{Z}^{-1}(\mu) \tilde{j}_0$, $\tilde{j}_0 = \bar{q}_0 \tilde{Q}_0$ can be considered similarly (results don't depend on its γ -matrix structure). Instead of (4), we have now

$$\tilde{C}(\mu) = \left(\frac{Z_q}{Z'_q} \right)^{1/2} \left(\frac{\tilde{Z}_Q}{\tilde{Z}'_Q} \right)^{1/2} \frac{\tilde{Z}'(\mu) \tilde{\Gamma}_0}{\tilde{Z}(\mu) \tilde{\Gamma}'_0}. \quad (12)$$

On-shell renormalization constant of the static quark field is $\tilde{Z}_Q = \left[1 - \left(d\tilde{\Sigma}/d\omega \right)_{\omega=0} \right]^{-1}$, where at two loops [4]

$$\left. \frac{d\tilde{\Sigma}}{d\omega} \right|_{\omega=0} = -C_F T_F \frac{g_0^4 M^{-4\varepsilon}}{(4\pi)^d} (d-1) I. \quad (13)$$

The bare vertex function at two loops is $\tilde{\Gamma}_0 = 1$ [4]. The ratio

$$\frac{\tilde{Z}}{\tilde{Z}'} = 1 + C_F T_F \left(\frac{\alpha_s}{4\pi\varepsilon} \right)^2 \left(1 - \frac{5}{6}\varepsilon \right) \quad (14)$$

can be obtained from [8]. Finally, we arrive at

$$\tilde{j}(M) = \left[1 + \frac{89}{36} C_F T_F \left(\frac{\alpha_s}{4\pi} \right)^2 \right] \tilde{j}'(M). \quad (15)$$

It was stated in [4] that it is impossible to match QCD heavy-light currents to HQET without heavy-quark loops by equating exclusive matrix elements. This statement is incorrect; it was based on eq. (2.1) of that paper, whose right-hand side should be multiplied by $Z_q^{1/2}$ in this case. Taking into account (15), we can write the matching coefficients for HQET without heavy-quark loops in the form (2.20) [4] with $a'_f = a_f + \frac{89}{36}$. Correspondingly, all numerical values of the coefficients of $(\alpha_s/\pi)^2$ in Table 1 [4] should be shifted by 0.103 in this case.

In conclusion, light-light quark currents should be adjusted according to (10) when crossing a heavy-flavour threshold; the heavy-light HQET current (say, $\bar{q}b$) should be adjusted according to (15) when crossing a threshold (say, of c quark).

I am grateful to K. G. Chetyrkin for useful discussions, and to T. Mannel for hospitality in Karlsruhe.

References

- [1] W. Wetzel, *Nucl. Phys.* **B196** (1982) 259;
W. Bernreuther and W. Wetzel, *Nucl. Phys.* **B197** (1982) 228;
W. Bernreuther, *Ann. Phys.* **1** (1983) 127; *Z. Phys.* **C20** (1983) 331.
- [2] S. A. Larin, T. van Ritbergen and J. A. M. Vermaseren, *Nucl. Phys.* **B438** (1995) 278.
- [3] K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, *Nucl. Phys.* **B510** (1998) 61.
- [4] D. J. Broadhurst and A. G. Grozin, *Phys. Rev.* **D52** (1995) 4082.
- [5] K. G. Chetyrkin, *Phys. Lett.* **B307** (1993) 169.
- [6] K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, *Nucl. Phys.* **B490** (1997) 19.
- [7] S. A. Larin and J. A. M. Vermaseren, *Phys. Lett.* **B259** (1991) 345;
S. A. Larin, in *Quarks-92*, ed. D. Yu. Grigoriev, V. A. Matveev, V. A. Rubakov and P. G. Tinyakov (World Scientific, Singapore, 1993), p. 201 (hep-ph/9302240);
Phys. Lett. **B303** (1993) 113.
- [8] D. J. Broadhurst and A. G. Grozin, *Phys. Lett.* **B267** (1991) 105.